



After your treatment

Your healthcare professional will review with you what to expect following treatment with JUVÉDERM™ injectable gel.

You should avoid the following for at least 24 hours:

- Strenuous exercise
- Consumption of alcoholic beverages
- Extended exposure to sun or heat

Exposure to any of the above may cause temporary redness, swelling, and/or itching at the injection sites. If you have swelling at the injection site, you can apply an ice pack for a brief period. Your healthcare professional will inform you when makeup may be applied after your treatment.

If you have any questions or concerns about your JUVÉDERM™ treatment, please contact your healthcare professional.



THE SCIENCE OF REJUVENATION™

©2007 Allergan, Inc. ™ mark owned by Allergan, Inc.

JUVÉDERM™ mark owned by Corneal Industrie SAS.

Re-order: M1656-01 7/07 701949



Your next
appointment

Physician name

Physician phone number

Patient name

Target date and time

JUVÉDERM™ Injectable Gel

Important Treatment Considerations for Patients

In the US, JUVÉDERM™ injectable gel is indicated for injection into the mid to deep dermis for correction of moderate to severe facial wrinkles and folds (such as nasolabial folds) and is generally well tolerated. The most commonly reported side effects are temporary injection site redness, swelling, pain/tenderness, firmness, lumps/bumps, and bruising. Exposure of the treated area to excessive sun and extreme cold weather should be minimized until any initial swelling and redness have resolved. If laser treatment, chemical peel, or any other procedure based on active dermal response is considered after treatment with JUVÉDERM™ injectable gel, there is a possible risk of an inflammatory reaction at the treatment site.

Patients who are using substances that can prolong bleeding, such as aspirin or ibuprofen, as with any injection, may experience increased bruising or bleeding at injection site. You should inform your physician before treatment if you are using these types of substances. As with all skin injection procedures, there is a risk of infection. JUVÉDERM™ injectable gel should be used with caution in patients on immunosuppressive therapy, or therapy used to decrease the body's immune response, as there may be an increased risk of infection. The safety of JUVÉDERM™ injectable gel in patients with a history of excessive scarring (eg, hypertrophic scarring and keloid formations) and pigmentation disorders has not been studied.

JUVÉDERM™ injectable gel should not be used in patients who have severe allergies marked by a history of anaphylaxis or history or presence of multiple severe allergies. JUVÉDERM™ injectable gel should not be used in patients with a history of allergies to Gram-positive bacterial proteins. The safety of JUVÉDERM™ injectable gel for use during pregnancy, in breastfeeding females, or in patients under 18 years has not been established. The safety and effectiveness of JUVÉDERM™ injectable gel for the treatment of areas other than facial wrinkles and folds (such as lips) have not been established in controlled clinical studies. For full risk information, please visit www.juvederm.com or call the Allergan Product Support line at 1-877-345-5372.

