



2680 S VAL VISTA DR STE 125 / GILBERT, AZ 85295
480.553.9909 WWW.COPPERCANYONDENTALAZ.COM

Extractions Post Operative Instructions

Avoid chewing for two hours following surgery, or until the numbness has completely worn off. Some discomfort after the extraction is normal.

The initial healing period usually takes one to two weeks, and you'll likely experience some swelling for the first forty-eight hours.

Be sure to take any antibiotic medication prescribed for you as it will help the area to heal. A pain medicine may also be recommended for your comfort and can be taken as directed.

You can also decrease pain and swelling by applying an ice pack twenty minutes on, twenty minutes off for the first six hours following the tooth extraction(s).

A blood clot will form on the extraction site, and this clot is vital to the healing process. To keep the clot intact, avoid touching the extraction site with your tongue or fingers, do not drink liquids through a straw, and do not spit or swish vigorously. Avoid blowing your nose or sneezing violently can also dislodge the blood clot and impair healing.

Do not rinse your mouth the day of the surgery. Smoking, or allowing food particles to pack into the tooth's socket, should be avoided, as both will significantly affect the healing process.

Twenty-four hours following the procedure, you can rinse gently with or a warm saltwater solution. (Dissolve one teaspoon of salt with one cup of warm water. Gently swish the solution around the affected area, and spit carefully.) You should do this two to three times each day for the week following the extraction.

Relax as much as possible and avoid all strenuous activities for the first twenty-four hours following surgery.



COPPER CANYON
D E N T A L

2680 S VAL VISTA DR STE 125 / GILBERT, AZ 85295
480.553.9909 WWW.COPPERCANYONDENTALAZ.COM

Once the numbness has worn off, you should eat, as nourishment is important to the healing process. Limit your diet to soft foods like yogurt, soft soups, ice cream, or soft-cooked eggs for the first forty-eight hours, and drink at least eight large glasses of water or fruit juice each day.

Keep your head elevated with pillows to control bleeding. We will give you a supply of gauze sponges to place over the bleeding area.

Change the pad as necessary, and use them until the bleeding stops completely. You can also bite gently but firmly on a moist tea bag for twenty minutes. Be sure to call our office if bleeding persists or increases.

The space left by the tooth will feel a bit strange to you at first. Eventually, new bone and gum tissue will grow into the gap left by the extraction.

Bleeding Control:

Slight bleeding or oozing after the surgery is normal and may occur at intervals for the first 24-48 hours. Forceful rinsing or excessive spitting can disrupt clotting, so do not rinse for at least 4 hours. If bleeding does persist:

- * Apply a moistened gauze pad, or tea bag, with moderate pressure to the bleeding site for 20 minutes. Repeat, if necessary.
- * Stay calm and quiet; lie down with the head elevated on 2-3 pillows.
- * If the bleeding persists do not hesitate to call our office.