



macedon family dentistry

joseph m. behrman, d.d.s. jeni l. behrman, d.m.d.

1212 route 31
p. o. box 862
macedon, new york 14502

phone (315) 986-3545
fax (315) 986-1074

INFORMATION FOLLOWING A TOOTH EXTRACTION

- **Bleeding** - After an extraction, a gauze pack is placed over the extraction site to prevent excessive bleeding and to promote the healing blood clot. Keep pressure on it for 30 - 45 minutes and replace if bleeding continues. Slight bleeding may occur up to 2 days. Avoid activities that could apply a suction action to the blood clot such as spitting or sucking through a straw. Avoid smoking for the next 24 hours, as it may cause the newly forming clot to be lost prematurely.
- **Rinsing** - Do not rinse your mouth today. Tomorrow you can rinse your mouth gently with a glass of warm water every 3-4 hours, especially after meals.
- **Swelling** - Following an extraction, some swelling and bruising may occur. A cold moist cloth or an ice bag applied to the cheek, will keep it to a minimum. Place on affected area for about 15 - 20 minutes of every hour for the next 6 hours.
- **Medications** - If over the counter pain medication does not relieve the discomfort you may experience, a stronger medication can be prescribed. Be sure to use all medications as directed.
- **Food** - A light diet with plenty of fluids is recommended the first day. Avoid carbonated or hot beverages. Chewing should be done away from the extraction site.
- **Oral Hygiene** - Continue brushing and flossing being extra gentle near the extraction site.
- **Chips** - During healing you may notice small bony fragments working their way through the gums. This is not a problem unless they become painful.